



TITLE:

# LEXICOGRAPHIC GROBNER BASES OF TORIC IDEALS ARISING FROM ROOT SYSTEMS (Algorithms in Algebraic Systems and Computation Theory)

AUTHOR(S):

Ohsugi, Hidefumi

---

CITATION:

Ohsugi, Hidefumi. LEXICOGRAPHIC GROBNER BASES OF TORIC IDEALS ARISING FROM ROOT SYSTEMS (Algorithms in Algebraic Systems and Computation Theory). 数理解析研究所講究録 2002, 1268: 73-76

ISSUE DATE:

2002-06

URL:

<http://hdl.handle.net/2433/42137>

RIGHT:

# LEXICOGRAPHIC GRÖBNER BASES OF TORIC IDEALS ARISING FROM ROOT SYSTEMS

大杉英史

HIDEFUMI OHSUGI

立教大学理学部数学科

Department of Mathematics, Rikkyo University

**ABSTRACT.** The present paper is a brief draft based on a joint work with Takayuki Hibi. Gröbner bases of toric ideals arising from root systems are studied.

## INTRODUCTION

Let  $\mathcal{A} \subset \mathbb{Z}^n$  be a finite set and let  $K[t, t^{-1}, s] = K[t_1, t_1^{-1}, \dots, t_n, t_n^{-1}, s]$  denote the Laurent polynomial ring over a field  $K$ . We associate each  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}^n$  with the monomial  $t^\alpha s = t_1^{\alpha_1} \dots t_n^{\alpha_n} s \in K[t, t^{-1}, s]$  and write  $\mathcal{R}_K[\mathcal{A}]$  for the subalgebra of  $K[t, t^{-1}, s]$  generated by all monomials  $t^\alpha s$  with  $\alpha \in \mathcal{A}$ . Let  $K[\mathbf{x}] = K[\{x_\alpha; \alpha \in \mathcal{A}\}]$  denote the polynomial ring in  $\#(\mathcal{A})$  variables over  $K$  and  $I_{\mathcal{A}} \subset K[\mathbf{x}]$  the kernel of the surjective homomorphism  $\pi : K[\mathbf{x}] \rightarrow \mathcal{R}_K[\mathcal{A}]$  defined by setting  $\pi(x_\alpha) = t^\alpha s$  for all  $\alpha \in \mathcal{A}$ . The ideal  $I_{\mathcal{A}}$  is called the *toric ideal* of the configuration  $\mathcal{A}$ . It is known [9] that if  $I_{\mathcal{A}}$  possesses a squarefree initial ideal, then the convex hull of  $\mathcal{A}$  possesses a unimodular triangulation.

Fix  $n \geq 2$ . Let  $\mathbf{e}_i$  denote the  $i$ -th unit coordinate vector of  $\mathbb{R}^n$ . We write  $\mathbf{A}_{n-1}^+$ ,  $\mathbf{B}_n^+$ ,  $\mathbf{C}_n^+$ ,  $\mathbf{D}_n^+$  and  $\mathbf{BC}_n^+$  for the set of positive roots of root systems  $\mathbf{A}_{n-1}$ ,  $\mathbf{B}_n$ ,  $\mathbf{C}_n$ ,  $\mathbf{D}_n$  and  $\mathbf{BC}_n$ , respectively ([3, pp. 64 – 65]):

$$\begin{aligned}\mathbf{A}_{n-1}^+ &= \{\mathbf{e}_i - \mathbf{e}_j; 1 \leq i < j \leq n\}; \\ \mathbf{B}_n^+ &= \{\mathbf{e}_i; 1 \leq i \leq n\} \cup \{\mathbf{e}_i + \mathbf{e}_j; 1 \leq i < j \leq n\} \cup \{\mathbf{e}_i - \mathbf{e}_j; 1 \leq i < j \leq n\}; \\ \mathbf{C}_n^+ &= \{2\mathbf{e}_i; 1 \leq i \leq n\} \cup \{\mathbf{e}_i + \mathbf{e}_j; 1 \leq i < j \leq n\} \cup \{\mathbf{e}_i - \mathbf{e}_j; 1 \leq i < j \leq n\}; \\ \mathbf{D}_n^+ &= \{\mathbf{e}_i + \mathbf{e}_j; 1 \leq i < j \leq n\} \cup \{\mathbf{e}_i - \mathbf{e}_j; 1 \leq i < j \leq n\}; \\ \mathbf{BC}_n^+ &= \mathbf{B}_n^+ \cup \mathbf{C}_n^+.\end{aligned}$$

Let, in addition,  $\tilde{\Phi}^+ = \Phi^+ \cup \{(0, 0, \dots, 0)\}$ , where  $\Phi = \mathbf{A}_{n-1}, \mathbf{B}_n, \mathbf{C}_n, \mathbf{D}_n$  or  $\mathbf{BC}_n$  and where  $(0, 0, \dots, 0)$  is the origin of  $\mathbb{R}^n$ .

In their combinatorial study of hypergeometric functions associated with root systems, Gelfand, Graev and Postnikov [2, Theorem 6.3] discovered a squarefree quadratic initial ideal of the toric ideal  $I_{\tilde{\mathbf{A}}_{n-1}^+}$  of  $\tilde{\mathbf{A}}_{n-1}^+$ . Moreover, for *any* subconfiguration  $\mathcal{A}$  of  $\mathbf{A}_{n-1}^+$ , the configuration  $\tilde{\mathcal{A}} = \mathcal{A} \cup (0, 0, \dots, 0)$  possesses a regular unimodular triangulation ([7, Example 2.4 (a)]). Stanley [8, Exercise 6.31 (b), p. 234] computed the Ehrhart polynomial of the convex polytope  $\text{conv}(\tilde{\mathbf{A}}_{n-1}^+)$ . Fong [1] constructed certain triangulations of the configurations  $\tilde{\mathbf{B}}_n^+ (= \text{conv}(\tilde{\mathbf{D}}_n^+) \cap \mathbb{Z}^n)$

and  $\text{conv}(\tilde{C}_n^+) \cap \mathbb{Z}^n (= \widetilde{BC}_n^+)$ , and computes the Ehrhart polynomials of  $\text{conv}(\tilde{B}_n^+)$  and  $\text{conv}(\tilde{C}_n^+)$ . The triangulations studied in [1] are, however, non-unimodular. Motivated by their results, Ohsugi–Hibi [6] showed that

**Proposition 0.1.** *Let  $\Phi \subset \mathbb{Z}^n$  be one of the root systems  $A_{n-1}$ ,  $B_n$ ,  $C_n$ ,  $D_n$  and  $BC_n$ . Then, there exists a reverse lexicographic order such that the initial ideal of  $I_{\Phi^+}$  is generated by squarefree quadratic monomials.*

Moreover, Ohsugi–Hibi [5] discussed subconfigurations  $\tilde{\mathcal{A}} = \mathcal{A} \cup \{(0, 0, \dots, 0)\}$  of  $\tilde{B}_n^+ \cup \tilde{C}_n^+$  which possesses a (regular) unimodular triangulation (i.e.,  $I_{\tilde{\mathcal{A}}}$  which possesses a squarefree initial ideal).

Hence, it is natural to study the same problem as above for  $I_{\Phi^+}$  where  $\Phi \subset \mathbb{Z}^n$  is one of the root systems  $A_{n-1}$ ,  $B_n$ ,  $C_n$ ,  $D_n$  and  $BC_n$ . (Then,  $I_{\Phi^+}$  is not generated by quadratic binomials if  $n \geq 6$ .)

### 1. SQUAREFREE LEXICOGRAPHIC INITIAL IDEALS

Let  $\Phi^+ \subset \mathbb{Z}^n$  denote one of the configurations  $A_{n-1}^+$ ,  $B_n^+$ ,  $C_n^+$ ,  $D_n^+$  and  $BC_n^+$ . Let  $K[A_{n-1}^+]$ ,  $K[B_n^+]$ ,  $K[C_n^+]$ ,  $K[D_n^+]$  and  $K[BC_n^+]$  denote the polynomial rings

$$\begin{aligned} K[A_{n-1}^+] &= K[\{f_{i,j}\}_{1 \leq i < j \leq n}], \\ K[B_n^+] &= K[\{y_i\}_{1 \leq i \leq n} \cup \{e_{i,j}\}_{1 \leq i < j \leq n} \cup \{f_{i,j}\}_{1 \leq i < j \leq n}], \\ K[C_n^+] &= K[\{a_i\}_{1 \leq i \leq n} \cup \{e_{i,j}\}_{1 \leq i < j \leq n} \cup \{f_{i,j}\}_{1 \leq i < j \leq n}], \\ K[D_n^+] &= K[\{e_{i,j}\}_{1 \leq i < j \leq n} \cup \{f_{i,j}\}_{1 \leq i < j \leq n}], \\ K[BC_n^+] &= K[\{a_i\}_{1 \leq i \leq n} \cup \{y_i\}_{1 \leq i \leq n} \cup \{e_{i,j}\}_{1 \leq i < j \leq n} \cup \{f_{i,j}\}_{1 \leq i < j \leq n}] \end{aligned}$$

over  $K$ . Write  $\pi : K[\Phi^+] \rightarrow K[t, t^{-1}, s]$  for the homomorphism defined by setting

$$\pi(a_i) = t_i^2 s, \quad \pi(y_i) = t_i s, \quad \pi(e_{i,j}) = t_i t_j s, \quad \pi(f_{i,j}) = t_i t_j^{-1} s.$$

Thus the kernel of  $\pi$  is the toric ideal  $I_{\Phi^+}$ .

First, an explicit initial ideals of  $I_{A_{n-1}^+}$  generated by squarefree monomials of degree  $\leq 3$  will be constructed. Let  $<_{lex}$  be the lexicographic order induced by the ordering of variables

$$f_{n-1,n} > f_{n-2,n-1} > f_{n-2,n} > \cdots > f_{1,2} > f_{1,3} > \cdots > f_{1,n},$$

and let  $<_{rev}$  be the reverse lexicographic order induced by the ordering of variables

$$f_{n-1,n} > f_{n-2,n} > f_{n-2,n-1} > \cdots > f_{2,3} > f_{1,n} > \cdots > f_{1,3} > f_{1,2}.$$

Then, the reduced Gröbner basis with respect to  $<_{lex}$  (and  $<_{rev}$ ) is as follows.

**Theorem 1.1** ([4]). *The set of the binomials*

$$\begin{aligned} f_{i,\ell} f_{j,k} - f_{i,k} f_{j,\ell}, & \quad i < j < k < \ell, \\ f_{i,j} f_{j,k} - f_{i,i+1} f_{i+1,k}, & \quad i+1 < j < k, \\ f_{i,j} f_{k,k+1} f_{k+1,\ell} - f_{i,i+1} f_{i+1,j} f_{k,\ell}, & \quad i+1 < j < k < \ell-1, \end{aligned}$$

is the reduced Gröbner basis of the toric ideal  $I_{A_{n-1}^+}$  with respect to both  $<_{lex}$  and  $<_{rev}$ , where the initial monomial of each binomial is the first monomial.

Then, we can associate the initial ideal of  $I_{A_{n-1}^+}$  with respect to  $<_{lex}$  with the regular unimodular triangulation  $\Delta_{<_{lex}}$ . A graph-theoretical characterization of the maximal faces of the triangulation  $\Delta_{<_{lex}}$  is given in [4].

Second, we discuss the existence of squarefree initial ideals of the toric ideal  $I_{\Phi^+}$  where  $\Phi \subset \mathbb{Z}^n$  is one of the root systems  $B_n$ ,  $C_n$ ,  $D_n$  and  $BC_n$ . The similar argument as in [5] plays an important role in the proof of Theorems 1.2 and 1.4.

Let  $<_{lex}^c$  be the lexicographic order induced by the ordering of variables

$$\begin{aligned} a_1 &> a_2 > \cdots > a_n \\ &> f_{n-1,n} > f_{n-2,n-1} > f_{n-2,n} > \cdots > f_{1,2} > f_{1,3} > \cdots > f_{1,n} \\ &> e_{n-1,n} > e_{n-2,n-1} > e_{n-2,n} > \cdots > e_{1,2} > e_{1,3} > \cdots > e_{1,n}. \end{aligned}$$

**Theorem 1.2.** *The initial ideal of the toric ideal  $I_{C_n^+}$  with respect to  $<_{lex}^c$  is generated by squarefree monomials.*

Let  $<_{lex}^d$  denote the lexicographic order obtained by restricting  $<_{lex}^c$  to  $K[\mathbf{D}_n^+]$ . By the elimination property of the lexicographic order  $<_{lex}^c$ , we have the following corollary from Theorem 1.2.

**Corollary 1.3.** *The initial ideal of the toric ideal  $I_{D_n^+}$  with respect to  $<_{lex}^d$  is generated by squarefree monomials.*

We now consider the root systems  $B_n$  and  $BC_n$ . Let  $<_{lex}^{bc}$  be the lexicographic order induced by the ordering of variables

$$\begin{aligned} a_1 &> a_2 > \cdots > a_n \\ &> e_{n-1,n} > e_{n-2,n-1} > e_{n-2,n} > \cdots > e_{1,2} > e_{1,3} > \cdots > e_{1,n} \\ &> y_1 > y_2 > \cdots > y_n \\ &> f_{n-1,n} > f_{n-2,n-1} > f_{n-2,n} > \cdots > f_{1,2} > f_{1,3} > \cdots > f_{1,n}. \end{aligned}$$

**Theorem 1.4.** *The initial ideal of the toric ideal  $I_{BC_n^+}$  with respect to  $<_{lex}^{bc}$  is generated by squarefree monomials.*

Let  $<_{lex}^b$  denote the lexicographic order obtained by restricting  $<_{lex}^{bc}$  to  $K[\mathbf{B}_n^+]$ . By the elimination property of the lexicographic order  $<_{lex}^{bc}$ , we have the following corollary from Theorem 1.4.

**Corollary 1.5.** *The initial ideal of the toric ideal  $I_{B_n^+}$  with respect to  $<_{lex}^b$  is generated by squarefree monomials.*

**Remark 1.6.** Let  $n \geq 6$  and let  $\Phi^+$  denote one of the configurations  $A_{n-1}^+$ ,  $B_n^+$ ,  $C_n^+$ ,  $D_n^+$  and  $BC_n^+$ . Then  $I_{\Phi^+}$  is not generated by quadratic binomials. Hence, in particular,  $I_{\Phi^+}$  does not possess a quadratic Gröbner basis.

## REFERENCES

- [1] W. Fong, Triangulations and Combinatorial Properties of Convex Polytopes, Dissertation, M. I. T., June, 2000.
- [2] I. M. Gelfand, M. I. Graev and A. Postnikov, Combinatorics of hypergeometric functions associated with positive roots, in "Arnold–Gelfand Mathematics Seminars, Geometry and Singularity Theory" (V. I. Arnold, I. M. Gelfand, M. Smirnov and V. S. Retakh, Eds.), Birkhäuser, Boston, 1997, pp. 205 – 221.
- [3] J. E. Humphreys, "Introduction to Lie Algebras and Representation Theory," Second Printing, Revised, Springer–Verlag, Berlin, Heidelberg, New York, 1972.
- [4] T. Kitamura, H. Ohsugi and T. Hibi, Gröbner bases associated with positive roots and Catalan numbers, preprint.
- [5] H. Ohsugi and T. Hibi, Unimodular triangulations and coverings of configurations arising from root systems, *J. Algebraic Combinatorics*, **14** (2001), 199 – 219.
- [6] H. Ohsugi and T. Hibi, Quadratic initial ideals of root systems, *Proc. Amer. Math. Soc.*, **130** (2002), 1913 – 1922.
- [7] R. P. Stanley, Decompositions of rational convex polytopes, *Ann. Discrete Math.* **6** (1980), 333 – 342.
- [8] R. P. Stanley, "Enumerative Combinatorics, Volume II," Cambridge University Press, Cambridge, New York, Sydney, 1999.
- [9] B. Sturmfels, "Gröbner Bases and Convex Polytopes," Amer. Math. Soc., Providence, RI, 1995.

Department of Mathematics,  
 Rikkyo University,  
 Nishi-Ikebukuro, Tokyo 171–8501, Japan  
 E-mail: ohsugi@rkmath.rikkyo.ac.jp